

NAVAL POSTGRADUATE SCHOOL

Monterey, California



**MODELING AND STATISTICAL ANALYSIS OF
BIOASSAY DATA: MEDAKA CELL PROLIFERATION
UNDER DEN AND TCE**

Donald P. Gaver
Patricia A. Jacobs

DTIC QUALITY INSPECTED 2

October 1994

Approved for public release; distribution is unlimited.

Prepared for:
U.S. Army Biomedical Research & Development Laboratory
Ft. Detrick, MD 21702-5010

19941227 060

NAVAL POSTGRADUATE SCHOOL
MONTEREY, CA 93943-5000


Rear Admiral T. A. Mercer
Superintendent

Harrison Shull
Provost

This report was prepared for and funded by the U.S. Army Biomedical Research and Development Laboratory, Ft. Detrick, MD.

Reproduction of all or part of this report is authorized.

This report was prepared by:



DONALD P. GAVAR, JR.
Professor of Operations Research



PATRICIA A. JACOBS
Professor of Operations Research

Reviewed by:

Released by:



PETER PURDUE
Professor and Chairman
Department of Operations Research



PAUL J. MARTO
Dean of Research

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE October 1994		3. REPORT TYPE AND DATES COVERED Technical
4. TITLE AND SUBTITLE Modeling and Statistical Analysis of Bioassay Data; Medaka Cell Proliferation under DEN and TCE			5. FUNDING NUMBERS RLLDG002JA	
6. AUTHOR(S) Donald P. Gaver and Patricia A. Jacobs				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Postgraduate School Monterey, CA 93943			8. PERFORMING ORGANIZATION REPORT NUMBER NPS-OR-94-014	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Biomedical Research & Development Laboratory Ft. Detrick, MD 21702-5010			10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES The views expressed in this report are those of the authors and do not reflect the official policy or position of the Department of Defense or the U.S. Government.				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) The response of <i>medaka</i> liver to the chemicals DEN and TCE is analyzed statistically. The analysis illustrates the application of methods useful in <i>environmetrics</i> , i.e. environmental statistics. It suggests an overall dose-response <i>effect</i> but not an easily-interpreted dose-response functional <i>relationship</i> .				
14. SUBJECT TERMS box plots, variance-stabilizing transformation, multiple comparisons, analysis of variance (ANOVA)			15. NUMBER OF PAGES 48	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT UL	

Modeling and Statistical Analysis of Bioassay Data: Medaka Cell Proliferation under DEN and TCE

by
Donald P. Gaver

and

Patricia A. Jacobs

**Department of Operations Research
Naval Postgraduate School
Monterey, CA 93943**

Accession For	
NTIS	CRA&I <input checked="" type="checkbox"/>
DTIC	TAB <input type="checkbox"/>
Unannounced <input type="checkbox"/>	
Justification _____	
By _____	
Distribution / _____	
Availability Codes	
Dist	Avail and/or Special
A-1	

1. INTRODUCTION

This report describes an analysis of cell proliferation data, by liver slice, from an experiment using Japanese Medaka. Previous work has used summary data from the same experiment (Gaver and Jacobs, 1994a,b). Another relevant reference is to Morris (1993). A brief description of the experiment follows.

The medaka are exposed to differing levels of DEN and TCE in tanks of water. The treatment groups are: control, 10 mg/ℓ DEN, 100 mg/ℓ DEN, 0.1 mg/ℓ TCE, (10 mg/ℓ DEN with 0.1 mg/ℓ TCE), 1 mg/ℓ TCE, and (10 mg/ℓ DEN with 1 mg/ℓ TCE). Each treatment group has two replicate tanks. Eight animals in each tank were sacrificed on 4 August, 1993; this is sacrifice B. Eight additional animals in each tank were sacrificed on 20 August 1993; this is sacrifice D.

Each sacrificed fish was exposed to BrdU for 72 hours prior to sacrifice; any cell that is in S-phase during this time has a BrdU marker. Each sacrificed fish is frozen and sliced longitudinally into 7-micron sections. A third of the slices are stained with another agent. This agent stains nuclei with the BrdU marker black; these nuclei are called *positive*. It is 5 of the latter stained slices that are analyzed subsequently.

Five slices containing a portion of the liver are considered for each fish. A region of interest (ROI) is marked on the slice; the ROI is chosen to attempt to maximize the number of hepatocytes and minimize the number of nonhepatocytes present. The area of all of the hepatocytes within the region of interest is measured, and the area of positive nuclei within the region of interest is measured. The number of hepatocytes in the ROI, and the number of positive hepatocytes in the ROI were also counted for half the fish in sacrifice B.

A count measure of cell proliferation, the *count index* (CI), for a slice is the number of positive hepatocytes in the ROI divided by the number of hepatocytes in the ROI, multiplied by 100. Evaluation of this measure is very labor intensive. As an alternative, the following *area index* (AI) is used

$$AI = \text{Area Index} = \frac{\text{Area of positive nuclei in the ROI}}{\text{Area of hepatocytes in the ROI}} \times 100.$$

The area index is easier to obtain; however, it does not quantify cells in S-phase exactly as does the CI since cells are of different size, as are the areas resulting from the slicing process, which are the result of a random intersection with the cell.

Figure 1 displays a plot of the count index (divided by 100) versus area index (divided by 100), computed by slice, for those fish of sacrifice B for which both measures are available, along with a simple unweighted least-squares-fitted

straight line; also displayed is the least-squares line equation with the standard errors of the coefficients displayed in parentheses below the coefficients. There appears to be a satisfactory linear relationship between the count index and area index, indicating that AI and CI are generally measuring the same response. However the variability of the area index increases as the count index increases; this increase is generally associated with high DEN and TCE dose levels; its biological interpretation is not yet available. It suggests that cell sizes become more variable under dosage.

Figure 2 is a display of the slice area indices (divided by 100), by fish, for the two control tanks. Note the variability between fish and the somewhat greater variability between fish in tank 2 as compared to those in tank 1.

Several slice area datum appear to be missing or are of doubtful validity. These have been deleted from analytical consideration. They are listed in Table 1. An alternative might have been to use robust statistical procedures throughout; such procedures automatically down-weight highly discrepant observations. Furthermore, examination of the weights indicates discrepancy so that explanations can be sought. It seems likely that robust methods should be more widely used in environmental toxicology. Robust statistical methods are discussed seriously in Cox and Hinkley (1974). A less advanced treatment appears in Koopmann (1987).

A summary of the findings of the data analysis is as follows.

- a. Available data from sacrifice B suggests that there is a reasonably strong linear association between the count index and the area index. For ease of analysis the area index has been used throughout.

TABLE 1
Slices Not Considered in the Data Analysis

MISSING SLICES: SACRIFICE B				
Missing Measurements are left blank.				
ID	No. of Slices	Tank	Treatment	Reason
9273802	All	8	0.1 TCE	Blank field
9273811	All	9	10 DEN, 0.1 TCE	Blank field

MISSING SLICES: SACRIFICE D				
Missing Values appear to be coded with 0.				
ID	No. of Slices	Tank	Treatment	Reason
9273986	All	2	Control	Area ROI = Positive Area = 0
9285049	All	10	1 TCE	Area ROI = Positive Area = 0
9274001	All	4	10 DEN	All positive areas = 0 with 2 slices having 1 mit. hep.
9285079	All	14	10 DEN, 1 TCE	All positive areas = 0 with 3 slices having 1 mit. hep.
9285022	1	7	0.1 TCE	Area of ROI = 247.16, others of order 7500
9285043	1	9	10 DEN, 0.1 TCE	Positive area of slice = 0; 0 mit. cells
9285073	1	13	10 DEN, 1 TCE	Positive area of slice = 0; 0 mit. cells

- b. There is evidence that the variances of the slice area indices over fish exposed to various treatments are approximately equal to the corresponding means. This relationship can lead to misleading conclusions if standard statistical procedures are used uncritically; furthermore, the results are less efficient than necessary. However, the variances of the *square roots* of the slice area

indices for fish subjected to the various treatment (DEN and TCE) levels appear approximately constant, i.e. far less dependent on the corresponding mean values. Consequently, the square roots of the area indices are used in subsequent data analyses. The underlying reason for the above data behavior is that positive counts are rare random events, hence tend to be approximately Poisson distributed. The square root transformation is known to stabilize the variance of such counts; see Miller (1986), p. 59. The same transformation should, and here does, stabilize the variance of count-associated areas.

- c. The means of the fish mean square root of area indices are considered for both sacrifices. There is generally more treatment effect for sacrifice B than for sacrifice D: examine p -values in lines 3 of Tables 2 and 3 for the overall analysis of variance indications to see that sacrifice D p -values are always larger than those for sacrifice B. Four out of five of the treatment means are significantly larger (at the 95% level) than that for the control for sacrifice B. There is no significant difference (95% level) between the treatment means and the control mean for sacrifice D. For sacrifice B, the treatment means for two out of the three levels of TCE with 10 mg/ ℓ DEN are significantly larger than those without DEN for sacrifice B; there is no significant difference for sacrifice D.
- d. The control mean for sacrifice D is significantly larger than the control mean for sacrifice B. The other treatment means for sacrifice D are not significantly different than those for sacrifice B.

- e. There is some suggestion that in the later sacrifice D, the presence of TCE *lowers* the mean of the area index. The biological mechanisms likely to explain this behavior are not yet available.

The above results are from one-way analysis of variance (ANOVA) of the square roots of the area indices, augmented by *multiple comparison* methods; the latter allow all possible pairwise comparisons to be made with a specified experiment-wise error rate, here 5%.

Brief Overall Summary of Findings to Date

The above can be briefly, and simplistically, summarized as follows. While it can be said that there is a statistically significant difference between mean responses ($\sqrt{\text{AI}}$) to the various treatments with DEN and TCE, no simple and interpretable dose-response patterns have been found. In particular, response does not appear to increase (or decrease) systematically with dose increase, where "dose" includes time of exposure as well as increases in chemical concentration dose levels. It remains to be seen whether the latter inconclusivity is lessened by the analysis of more data (later sacrifices), by finding that experimental problems or biases occurred, *or*, more exciting, that the dose-responses observed can be explained by biological mechanism, and that the findings essentially reappear when further experiments and data analyses are conducted.

Section 2 presents results of graphical displays of the data. Section 3 presents results of exploratory analyses of variance. Results of exploratory linear regression models are presented in Section 4. Multiple comparison results appear in Section 5.

2. GRAPHICAL SUMMARIES OF THE SLICE AREA INDICES

Figure 3a (respectively 3b) displays boxplots of the slice area indices by tank for sacrifice B, (respectively sacrifice D). The boxplots may be viewed as a graphical one-way analysis of variance. There is some tank effect within a treatment group. The greatest dose-response effect is clearly for the 100 DEN treatment.

Figures 4a and 4b display plots of the mean of the area indices for each tank versus the variance of the area indices for each tank. Also displayed is a 45° line. There appears to be a linear relationship between the mean and variance; in fact, the variances appear to be approximately equal to the means. This relationship between the means and variances may lead to misleading results if analysis of variance techniques are applied directly to the slice area indices; Miller (1986) and Box (1954) discuss the effects of inequality of variance on one-way analysis of variance. It is believed that such effects may well appear often in environmental toxicology, particularly where *counts*, or count-like phenomena, are found.

As noted earlier, one standard transformation that can be applied to data with variances approximately equal to means to attempt to make the variance of the transformed data more nearly constant is the *square root transformation*; see Miller (1986). Figures 5a and 5b display plots of the mean of the square root of the area indices for each tank versus the variance of the square root of the area indices for each tank. The variances now appear unrelated to the corresponding means. Figures 6a and 6b display boxplots of the square roots of the slice area indices by tank. Note that the lengths (heights between quartiles) of the boxes are less variable than are those for the boxplots of the raw area indices themselves. In the remainder of this paper the square root of area indices will be used.

3. EXPLORATORY ANALYSES OF VARIANCE

Results of exploratory analyses of variances appear in Tables 2 – 4. The basic data are summaries of area indices and the square roots of area indices for each fish. Since the boxplots indicate that the 100 DEN treatment is associated with much larger area indices, the analyses of variances were done with and without 100 DEN. Table 2 shows the results by tank. Recall that small p -values will indicate tank, hence treatment, effect. The first row of the table indicates that the tank means of the fish mean area index are significantly different. However the second row of the table indicates that the tank means of the logarithms of the variance (log variance) of area indices for each fish are also significantly different. The graphical analysis has already indicated this difference. Now recall that one of the assumptions of analysis of variance is that the data come from populations with equal variance, so apply the square-root transformation. Rows 3 and 4 report results using the square roots of the area indices. Note that the results of row 4 indicate that there is no significant difference between the tank means of the log variance of the slice square root of area index for each fish. However, the results of row 3 indicate that there *is* significant difference between the tank *means* of the mean of the slice square root area indices for each fish; note that the p -values for the tank means are smaller using the fish mean square root of area indices rather than the raw (untransformed) area indices for the analysis of variance (without the 100 DEN treatment). Thus, the difference in variances of the raw (untransformed) area indices appears to have masked some of the difference in means, presumably resulting from treatment effects.

Table 3 displays results for analyses of variance, with the two tanks in each treatment combined. Once again, the analysis of variance using the mean slice

square root of the area indices for each fish indicates that there is a significant difference between treatment means of the means, even without 100 DEN.

TABLE 2
***p*-Values for Exploratory ANOVA of Area Indices (AI) by Tank**
 (AI = [(Positive Area)/ROI Area] × 100)
 (Small *p*-Values Indicate Chemical-Tank Effect)

	Sacrifice B		Sacrifice D	
Data	with 100 DEN	without 100 DEN	with 100 DEN	without 100 DEN
Mean slice AI for each fish in a tank	$< 10^{-16}$	3×10^{-5}	6×10^{-16}	2.9×10^{-2}
Log variance of slice AI for each fish in a tank	4×10^{-11}	2×10^{-5}	3×10^{-2}	0.54
Mean slice $\sqrt{\text{AI}}$ for each fish in a tank	4×10^{-16}	8×10^{-7}	1×10^{-14}	1.6×10^{-2}
Log variance of slice $\sqrt{\text{AI}}$ for each fish in a tank	0.16	0.25	0.89	0.97

TABLE 3
***p*-Values for Exploratory ANOVA of Area Indices (AI) by Treatment**
 (AI = [(Positive Area)/ROI Area] × 100)
 Treatment: 2 Tanks Combined
 (Small *p*-Values Indicate Treatment Effect)

	Sacrifice B		Sacrifice D	
Data	with 100 DEN	without 100 DEN	with 100 DEN	without 100 DEN
Mean slice AI for each fish in a treatment	1×10^{-16}	1×10^{-5}	1×10^{-16}	2×10^{-2}
Log variance of slice AI for each fish in a treatment	4×10^{-13}	3×10^{-6}	1×10^{-4}	0.33
Mean slice $\sqrt{\text{AI}}$ for each fish in a treatment	2×10^{-16}	9×10^{-7}	4×10^{-16}	9×10^{-3}
Log variance of slice $\sqrt{\text{AI}}$ for each fish in a treatment	0.19	0.46	0.56	0.89

Table 4 displays results of an analysis of variance of the mean square root of area indices for each fish in a treatment but without the control. Once again there is evidence of significant differences between the treatment means.

We conclude that there is a definite treatment effect, i.e. response to different treatment levels, even if no treatment = control and the “strong” 100 DEN treatment responses are removed.

TABLE 4
***p*-Values for Exploratory ANOVA of Area Indices (AI) by Treatment**
Without Control
(AI = [(Positive Area)/ROI Area] × 100)
Treatment: 2 Tanks Combined
(Small *p*-Values Indicate Treatment Effect)

	Sacrifice B		Sacrifice D	
Data	with 100 DEN	without 100 DEN	with 100 DEN	without 100 DEN
Mean $\sqrt{\text{AI}}$ for fish	0	1.9×10^{-3}	5.9×10^{-15}	7.4×10^{-3}

4. EXPLORATORY LINEAR REGRESSION

Tables 5 – 8 report results of fitting exploratory linear regression models to the square root of the slice area indices for each fish. For Tables 5 and 6, the covariates are the level of DEN minus its mean; the level of TCE minus its mean; and an interaction term: {(level of DEN minus its mean) times (level of TCE minus its mean)}. The means were subtracted to give the interaction term a value other than 0 if the level of DEN or the level of TCE is 0. The linear regressions were fit with and without the 100 DEN treatment. The results of Table 5 are for sacrifice B and those of Table 6 are for sacrifice D.

TABLE 5
Sacrifice B: \sqrt{AI}
Linear Regression Coefficient Estimates with Standard Error
and 95% Normal Confidence Intervals

<i>WITHOUT 100 DEN</i>					
CONSTANT	DEN- \overline{DEN}	TCE- \overline{TCE}	(DEN- \overline{DEN}) \times (TCE- \overline{TCE})	R ²	s.e.
(SE) [CI]	(SE) [CI]	(SE) [CI]	(SE) [CI]	— —	— —
1.88	0.037	0.526	0.025	0.15	0.50
(0.067)	(0.005)	(0.148)	(0.010)		
[1.75,2.01]	[0.028,0.046]	[0.237,0.816]	[0.005,0.045]		
$\overline{DEN} = 5.0$ $\overline{TCE} = 0.372$					
<i>WITH 100 DEN</i>					
1.77	0.029	0.710	0.039	0.56	0.50
(0.040)	(0.003)	(0.117)	(0.008)		
[1.69,1.85]	[0.024,0.033]	[0.48,0.94]	[0.02,0.05]		
$\overline{DEN} = 18.81$ $\overline{TCE} = 0.318$					

TABLE 6
Sacrifice D: \sqrt{AI}
Linear Regression Coefficient Estimates with Standard Error
and 95% Normal Confidence Intervals

<i>WITHOUT 100 DEN</i>					
CONSTANT	DEN- \overline{DEN}	TCE- \overline{TCE}	(DEN- \overline{DEN}) \times (TCE- \overline{TCE})	R ²	s.e.
(SE) [CI]	(SE) [CI]	(SE) [CI]	(SE) [CI]	— —	— —
1.89	0.028	-0.402	-0.025	0.08	0.50
(0.07)	(0.005)	(0.151)	(0.01)		
[1.76,2.03]	[0.02,0.04]	[-0.70,-0.11]	[-0.05,-0.005]		
$\overline{DEN} = 4.88$ $\overline{TCE} = 0.37$					
<i>WITH 100 DEN</i>					
1.68	0.013	-0.050	0.0005	0.45	0.50
(0.04)	(0.003)	(0.12)	(0.008)		
[1.60,1.76]	[0.008,0.018]	[-0.29,0.19]	[-0.02,0.02]		
$\overline{DEN} = 19.05$ $\overline{TCE} = 0.315$					

In both tables the values of R^2 are small when the 100 DEN data are excluded. This implies that a linear function of the above explanatory variables does not explain the data well. However, the standard errors of the estimates are also small. This behavior suggests that, although there may be an association between levels of DEN and TCE and the square root of the area index, that association is not linear. Note that for sacrifice B all of the estimates of the coefficients are significantly positive, suggesting that increasing levels of DEN and TCE are associated with higher area indices. However, for sacrifice D, the estimate of the coefficient of the level of TCE and the estimate of the coefficient of the interaction term coefficient are significantly negative for the regression, even without the 100 DEN treatment. This suggests that for the later sacrifice, exposure to TCE may have an inhibitory effect. The only estimate of covariate that is significantly different from 0 for sacrifice D when 100 DEN is included is exposure to DEN.

Tables 7 and 8 report results of fitting linear regressions with the covariate being the level of TCE exposure to data from fish not exposed to DEN and to data from fish exposed to 10 mg/ ℓ DEN; the dependent variable is the square root of the slice area index. The R^2 values are very small, indicating a lack of linear fit, but the estimate coefficients for TCE in the regressions using data from fish exposed to 10 mg/ ℓ DEN are formally significant. Once again, this behavior suggests that there may be an association but the association is not linear.

For sacrifice B, there was no significant effect for the level of TCE for the fish not exposed to DEN; for those fish exposed to 10 mg/ ℓ DEN the coefficient for level of TCE is significantly positive indicating that increasing levels of TCE are associated with increasing (square roots of) area indices.

TABLE 7
Sacrifice B: \sqrt{AI}
Linear Regression Coefficient Estimates with Standard Error
and 95% Normal Confidence Intervals
(Replicate Tanks Pooled)

<i>NO DEN</i>				
CONSTANT (SE) [CI]	TCE (SE) [CI]	DEN (SE) [CI]	R ² — —	s.e. — —
1.18	0.056	—	0.002	0.55
(0.047)	(0.080)	—		
[1.09, 1.27]	[-0.10, 0.21]	—		
<i>10 DEN</i>				
1.47	0.31	—	0.09	0.44
(0.037)	(0.06)	—		
[1.40, 1.54]	[0.18, 0.43]	—		
<i>0 DEN and 10 DEN</i>				
1.14	0.18	0.038	0.14	0.50
(0.038)	(0.051)	(0.005)		
[1.06, 1.21]	[0.08, 0.28]	[0.29, 0.047]		

In sacrifice D, for those fish not exposed to DEN, the estimate of the coefficient of TCE is not significantly different than 0. However, for those fish exposed to 10 mg/ ℓ DEN the estimate of the coefficient of TCE is significantly *negative*, suggesting that for the fish of the later sacrifice that were exposed to DEN, the *greater* the level of TCE exposure, the *smaller* the (square root of) the area index. This effect calls for biological explanation.

TABLE 8

Sacrifice D: \sqrt{AI}
 Linear Regression Coefficient Estimates with Standard Error
 and 95% Normal Confidence Intervals
 (Replicate Tanks Pooled)

<i>NO DEN</i>				
CONSTANT (SE) [CI]	TCE (SE) [CI]	DEN (SE) [CI]	R ² — —	s.e. — —
1.35	0.059	—	0.004	0.43
(0.037)	(0.062)	—		
[1.27, 1.42]	[-0.063, 0.181]	—		
<i>10 DEN</i>				
1.71	-0.189	—	0.02	0.56
(0.048)	(0.083)	—		
[1.61, 1.80]	[-0.352, -0.026]	—		
<i>0 DEN and 10 DEN</i>				
1.39	-0.061	0.027	0.07	0.50
(0.038)	(0.052)	(0.005)		
[1.32, 1.47]	[-0.16, 0.04]	[0.017, 0.036]		

5. MULTIPLE COMPARISONS

The exploratory analyses of variances strongly rejected the null hypothesis that all the treatment means (even without the 100 DEN treatment) of the fish mean square root of the area indices are equal. Rejection of the null hypothesis does not indicate specifically which means are not equal. A method for discovering which means differ is called a multiple comparisons procedure. There are a number of different *multiple comparisons* procedures in the literature; see Miller (1981). We will use two of them.

5.1 Simultaneous Confidence Intervals using Studentized Range Distribution

The first procedure uses the studentized range distribution to construct simultaneous confidence statements about the true values of all differences of the treatment means. Table 9 describes the procedure to obtain simultaneous 95% confidence intervals for all differences of treatment means for one sacrifice without the 100 DEN treatment. The original procedure requires that there be an equal number of fish in each treatment. However, Ott *et al.* suggest step 4 in Table 9 if the number of fish in each treatment do not differ by much.

5.1a Treatment Means Minus Control Mean

Figure 7 presents some of the 95% simultaneous confidence intervals of the differences of treatment means for sacrifice B. It shows the confidence intervals for the treatment means minus the control mean for the fish mean square root of the area indices. Note that 4 out of the 5 intervals are significantly above 0 indicating that the treatments are associated with a larger mean square root area indices than those for the control. The greatest difference is that for the treatment of 10 mg/ℓ DEN with 1 mg/ℓ TCE.

Figure 8 presents some of the 95% simultaneous confidence intervals for the later sacrifice D. It shows the confidence intervals for the treatment mean minus the control mean. None of the treatment means is significantly different from the control mean.

5.1b Treatment Means for Treatments with Exposure to 10 mg/ℓ DEN Minus Those without Exposure to 10 mg/ℓ DEN

Figure 9 displays the sacrifice B 95% simultaneous confidence intervals for the difference between the treatment means for mean fish square root of the area indices for those treatments *with* 10 mg/ℓ DEN minus the treatment means for those treatments *without* 10 mg/ℓ DEN, by level of TCE exposure. Note that the

treatment means with 10 mg/ℓ DEN is significantly larger than that without for 0 mg/ℓ TCE and 1 mg/ℓ TCE. There is no significant difference for 0.1 mg/ℓ TCE.

TABLE 9

To Obtain Tukey (Studentized Range Distribution) Simultaneous 95% Confidence Intervals for Treatment Mean Differences, One Sacrifice

1.	There are 6 treatments: Control; 10 DEN; 0.1 TCE; 10 DEN with 0.1 TCE; 1 TCE; and 10 DEN with 1 TCE.
2.	There are ~88 within degrees of freedom $\left(\sum_{i=1}^6 (n_i - 1) \right)$ where n_i is the number of fish in treatment i .
3.	The 0.05 percentage point for the studentized range for 60 within-degrees-of-freedom and 6 treatment means is 4.16, from published tables, <i>BIOMETRIKA Tables for Statisticians</i> , Vol. 1. This is larger than the percentage point for 88 within-degrees-of-freedom. Thus, the constructed confidence intervals will be conservative: one can truly say that <i>all</i> pairwise difference comparisons are made with (95%) confidence.
4.	Since the number of fish per treatment differs somewhat (due to missing fish) the harmonic mean of the number of fish per treatment is used $n = \frac{6}{\frac{1}{n_1} + \dots + \frac{1}{n_6}}$ where n_i is the number of fish in treatment i .
5.	The mean square within is $\frac{\sum_{i=1}^6 \sum_j (y_{ij} - \bar{y}_i)^2}{\sum_{i=1}^6 (n_i - 1)} = \text{MS}(\text{within})$ where y_{ij} is the mean $\sqrt{\text{AI}}$ for fish j in treatment i and \bar{y}_i is the mean of the mean $\sqrt{\text{AI}}$ for the fish in treatment i .
6.	95% confidence intervals for all pairs of means μ_i and $\mu_{i'}$ $(\bar{y}_i - \bar{y}_{i'}) \pm (4.16) \sqrt{\text{MS}(\text{within})/n}$

Figure 10 displays a similar plot for the later sacrifice D. There is no significant difference between the treatment means with 10 mg/ℓ DEN and those without.

5.1c Treatment Means for Sacrifice D Minus Treatment Means for Sacrifice B

Simultaneous confidence intervals are computed for all differences of the treatment means of the fish mean square root of the area indices for sacrifices B and D combined. Figure 11 displays six of the 95% simultaneous confidence intervals. It displays the 95% confidence intervals for the difference in treatment means between sacrifice D and sacrifice B. The only significant difference is for the control where the mean of the fish mean square root of area indices for sacrifice D is significantly larger than that for sacrifice B.

5.2 Studentized Maximum Modulus Confidence Intervals

Simultaneous confidence intervals for the treatment means themselves can be constructed using the studentized maximum modulus procedure; cf. Miller (1981). The procedure is as follows for the mean fish square root area index for the three treatment groups (10 mg/ℓ DEN, 0 mg/ℓ TCE), (10 mg/ℓ DEN, 0.1 mg/ℓ TCE), and (10 mg/ℓ DEN, 1 mg/ℓ TCE) for 1 sacrifice.

To Obtain Simultaneous Confidence Intervals for 3 Treatment Means using the Studentized Maximum Modulus Distribution

1. Compute the within degrees of freedom for the three treatments.

$$d = \sum_{i=1}^3 (n_i - 1)$$

where n_i is the number of fish in treatment i .

2. Compute the mean square within

$$MS(\text{within}) = \frac{\sum_{i=1}^3 \sum_j (y_{ij} - \bar{y}_{i.})^2}{\sum_{i=1}^3 (n_i - 1)}$$

where y_{ij} is the mean square root of the area indices for fish j in treatment i and $\bar{y}_{i.}$ is the mean of the fish means in treatment i .

3. Find the upper 0.05 point of the studentized maximum modulus distribution with parameters 3 (treatments) and d degrees of freedom, $m(3, d)$. Tables can be found in Miller (1981).
4. The three simultaneous 95% confidence intervals are

$$\bar{y}_{i.} \pm m(3, d) \sqrt{MS(\text{within})/n_i}.$$

Figure 12 displays the 95% simultaneous confidence intervals for sacrifice B for the means of the fish mean of the square root of the slice area indices for those treatments having fish exposed to 10 mg/ℓ DEN. The means appear about the same for 0 mg/ℓ TCE and 0.1 mg/ℓ TCE. The mean for 1 mg/ℓ TCE appears to be somewhat larger.

Figure 13 displays the 95% simultaneous confidence intervals for sacrifice D for the means of those treatments with 10 mg/ℓ DEN by level of TCE. There is some suggestion that the presence of TCE is associated with a *lower* mean of the fish mean square root of the slice area indices.

CONCLUSION

The above analyses illustrate the use of statistical methods appropriate for the kinds of data obtained by the medaka experiments. The methods of


transformation, analysis of variance, and multiple comparisons are useful and powerful for the initial data analyses, suggesting some surprising dose-response relations that are worthy of careful further biological investigation and explanation. Alternative methods can also be applied, and should yield the same general insights.

REFERENCES


- Bickel, P. J. and Doksum, K. A. *Mathematical Statistics: Basic Ideas and Selected Topics*, Holden-Day, Inc., San Francisco, 1977.
- Box, G. E. P. "Some theorems on quadratic forms applied in the study of analysis of variance problems, I. Effect of inequality of variance in the one-way classification." *Annals of Mathematical Statistics*, **25**, pp. 290-302.
- Cox, D. R. and Hinkley, D. V. *Theoretical Statistics*, Chapman and Hall, New York, 1974.
- Gaver, D. P. and Jacobs, P. A. "Assessment of liver modification and cell proliferation in medaka under DEN and TCE dosage, using data available 4/26/94, suggestions for statistical analysis and results from exploratory data analysis." Working Paper, April 26, 1994.
- Gaver, D. P. and Jacobs, P. A. "Comparison of area indices in medaka livers for sacrifices at different dose time combinations using data available 5/19/94." Working Paper, May 19, 1994.
- Huff, J. "Absence of morphologic correlation between chemical toxicity and chemical carcinogenesis." *Environmental Health Perspectives*, **101** (Suppl. 5) pp. 45-54, 1993.
- Koopmann, L. H. *Introduction to Contemporary Statistical Methods*, Duxbury Press, Boston, 1987.
- Miller, R. G. Jr. *Beyond ANOVA, Basics of Applied Statistics*, John Wiley & Sons, New York, 1986.
- Miller, R. G. Jr. *Simultaneous Statistical Inference*, Second Edition, Springer-Verlag, New York, 1981.
- Morris, R. W. "Analysis of cell proliferation data." *Environmental Health Perspectives*, **101** (Suppl. 5) pp. 73-78, 1993.
- Ott, L. and Hildebrand, D. K. *Statistical Thinking for Managers*, PWS Publishers, Boston, 1983.

AREA INDEX VERSUS COUNT INDEX BY SLICE

SCATTER PLOT, SSZ=275



RATIO POSITIVE AREA TO ROI AREA-TANK 1



RATIO POSITIVE AREA TO ROI AREA--TANK 2




Figure 2

SACRIFICE B: AREA INDICES

$((\text{POSITIVE AREA}) \div (\text{AREA ROI})) \times 100$ BY SLICE FROM ALL FISH




Figure 3a

SACRIFICE D: AREA INDICES

$((\text{POSITIVE AREA}) \div (\text{AREA ROI})) \times 100$ BY SLICE FROM ALL FISH





Figure 3b
24


SACRIFICE B: PLOT MEAN VS VARIANCE FOR EACH TANK

VARIANCE OF RATIO OF (POSITIVE AREA ÷ AREA ROI) × 100
USING SLICES WITHIN A TANK



SACRIFICE D: PLOT MEAN VS VARIANCE FOR EACH TANK

VARIANCE OF RATIO OF (POSITIVE AREA ÷ AREA ROI) × 100
USING SLICES WITHIN A TANK



SACRIFICE B: SQUARE ROOT AREA INDEX BY TANK




Figure 5a
27

SACRIFICE D: SQUARE ROOT AREA INDEX BY TANK





Figure 5b
28


SACRIFICE B

SQUARE ROOT AREA INDICES $((\text{POSITIVE AREA}) \div (\text{AREA ROI})) \times 100$
BY SLICE FROM ALL FISH



SACRIFICE D

SQUARE ROOT AREA INDICES ((POSITIVE AREA) ÷ (AREA ROI)) × 100
BY SLICE FROM ALL FISH



SACRIFICE B: SQUARE ROOT OF AREA INDICES

MEAN OF TREATMENT MEANS—MEAN OF CONTROL MEANS
+/- 95 PERC TUKEY SIMUL CONF LIMITS FOR ALL DIFF






Figure 7
31

SACRIFICE D: SQUARE ROOT OF AREA INDICES


MEAN OF TREATMENT MEANS—MEAN OF CONTROL MEANS
+/- 95 PERC TUKEY SIMUL CONF LIMITS FOR ALL DIFF



SACRIFICE B:MEAN(10 DEN)-MEAN(ODEN)
 95 PERCENT STUDENTIZED RANGE SIMULTANEOUS CONFIDENCE INTERVALS
 FOR ALL TREATMENT MEAN DIFFERENCES



SACRIFICE D:MEAN(10 DEN)–MEAN(ODEN)
 95 PERCENT STUDENTIZED RANGE SIMULTANEOUS CONFIDENCE INTERVALS
 FOR ALL TREATMENT MEAN DIFFERENCES



SQUARE ROOT OF AREA INDICES

MEAN OF MEANS BY TREATMENT FOR SACR D-SACR B
95 PERCENT CONF LIMITS OVER ALL DIFF




Figure 11

SACRIFICE B:10 DEN; SQUARE ROOT OF AREA INDICES
 STUDENTIZED MAXIMUM MODULUS SIMULTANEOUS CONFIDENCE INTERVALS
 95 PERCENT




Figure 12
 36

SACRIFICE D:10 DEN; SQUARE ROOT OF AREA INDICES
STUDENTIZED MAXIMUM MODULUS SIMULTANEOUS CONFIDENCE INTERVALS
95 PERCENT




Figure 13

INITIAL DISTRIBUTION LIST

1. Research Office (Code 08)1
 Naval Postgraduate School
 Monterey, CA 93943-5000

2. Dudley Knox Library (Code 52)2
 Naval Postgraduate School
 Monterey, CA 93943-5002

3. Defense Technical Information Center.....2
 Cameron Station
 Alexandria, VA 22314

4. Department of Operations Research (Code OR).....1
 Naval Postgraduate School
 Monterey, CA 93943-5000

5. Prof. Donald P. Gaver (Code OR/Gv).....5
 Naval Postgraduate School
 Monterey, CA 93943-5000

6. Prof. Patricia A. Jacobs (Code OR/Jc).....5
 Naval Postgraduate School
 Monterey, CA 93943-5000

7. Dr. J. Abrahams, Code 1111, Room 6071
 Mathematical Sciences Division, Office of Naval Research
 800 North Quincy Street
 Arlington, VA 22217-5000

8. Dr. John Bailar1
 468 N St. NW
 Washington, DC 20024

9. Dr. David Brillinger.....1
 Statistics Department
 University of California
 Berkeley, CA 94720

10. Dr. David Burman.....1
AT&T Bell Telephone Laboratories
600 Mountain Avenue
Murray Hill, NJ 07974

11. Prof. Brad Carlin.....1
School of Public Health
University of Minnesota
Mayo Bldg. A460
Minneapolis, MN 55455

12. Dr. Robert Carpenter1
NAMRI/Navy Toxicology Detachment
Bldg. 433, Area B
Wright-Patterson AFB, OH 45433-6503

13. Center for Naval Analyses.....1
4401 Ford Avenue
Alexandria, VA 22302-0268

14. Prof. H. Chernoff.....1
Department of Statistics
Harvard University
1 Oxford Street
Cambridge, MA 02138

15. Dr. Edward G. Coffman, Jr.....1
AT&T Bell Telephone Laboratories
600 Mountain Avenue
Murray Hill, NJ 07974

16. Dr. John Copas.....1
Dept. of Mathematics,
University of Birmingham
P. O. Box 363
Birmingham B15 2TT
ENGLAND

17. Professor Sir David Cox.....1
Nuffield College
Oxford, OXI INF
ENGLAND

18. Professor H. G. Daellenbach.....1
Department of Operations Research
University of Canterbury
Christchurch, NEW ZEALAND
19. Dr. S. R. Dalal.....1
Bellcore
445 South Street
Morristown, NJ 07962-1910
20. Dr. D. F. Daley1
Statistic Dept. (I.A.S.)
Australian National University
Canberra, A.C.T. 2606
AUSTRALIA
21. Dr. B. Doshi1
AT&T Bell Laboratories
HO 3M-335
Holmdel, NJ 07733
22. Naihua Duan.....1
RAND Corporation
Santa Monica, CA 90406
23. Prof. Bradley Efron.....1
Statistics Dept.
Sequoia Hall
Stanford University
Stanford, CA 94305
24. Dr. Guy Fayolle.....1
I.N.R.I.A.
Dom de Voluceau-Rocquencourt
78150 Le Chesnay Cedex
FRANCE
25. Dr. M. J. Fischer5
Defense Communications Agency
1860 Wiehle Avenue
Reston, VA 22070

26. Prof. George S. Fishman.....1
 Curr. in OR & Systems Analysis
 University of North Carolina
 Chapel Hill, NC 20742

27. Henry S. Gardner1
 U.S. Army Biological Research & Development Laboratory
 Ft. Detrick, MD 21702-5010

28. Dr. Neil Gerr1
 Office of Naval Research
 Arlington, VA 22217

29. Prof. Linda V. Green.....1
 Graduate School of Business
 Columbia University
 New York, NY 10027

30. Dr. D. C. Hoaglin1
 Department of Statistics
 Harvard University
 1 Oxford Street
 Cambridge, MA 02138

31. Prof. D. L. Iglehart.....1
 Dept. of Operations Research
 Stanford University
 Stanford, CA 94350

32. Institute for Defense Analysis.....1
 1800 North Beauregard
 Alexandria, VA 22311

33. Prof. J. B. Kadane.....1
 Dept. of Statistics
 Carnegie-Mellon University
 Pittsburgh, PA 15213

34. Dr. Jon Kettenring.....1
 Bellcore
 445 South Street
 Morris Township, NJ 07962-1910

35. Koh Peng Kong.....1
 OA Branch, DSO
 Ministry of Defense
 Blk 29 Middlesex Road
 SINGAPORE 1024

36. Dr. Prabha Kumar.....1
 Defense Communications Agency.....1
 1860 Wiehle Avenue
 Reston, VA 22070

37. Dr. A. J. Lawrence.....1
 Dept. of Mathematics,
 University of Birmingham
 P. O. Box 363
 Birmingham B15 2TT
 ENGLAND

38. Prof. M. Leadbetter.....1
 Department of Statistics
 University of North Carolina
 Chapel Hill, NC 27514

39. Prof. J. Lehoczky1
 Department of Statistics
 Carnegie-Mellon University
 Pittsburgh, PA 15213

40. Dr. Colin Mallows.....1
 AT&T Bell Telephone Laboratories
 600 Mountain Avenue
 Murray Hill, NJ 07974

41. Prof. R. Douglas Martin.....1
 Department of Statistics, GN-22
 University of Washington
 Seattle, WA 98195

42. Prof. M. Mazumdar1
 Dept. of Industrial Engineering
 University of Pittsburgh
 Pittsburgh, PA 15235

43. Dr. Sati Mazumdar1
 Biostatistics Dept.
 University of Pittsburgh
 Graduate School of Public Health
 Pittsburgh, PA 15261

44. Prof. Carl N. Morris.....1
 Statistics Department
 Harvard University
 1 Oxford St.
 Cambridge, MA 02138

45. Prof. F. W. Mosteller1
 Department of Statistics
 Harvard University
 1 Oxford St.
 Cambridge, MA 02138

46. Operations Research Center, Rm E40-164.....1
 Massachusetts Institute of Technology
 Attn: R. C. Larson and J. F. Shapiro
 Cambridge, MA 02139

47. Dr. John Orav.....1
 Biostatistics Department
 Harvard School of Public Health
 677 Huntington Ave.
 Boston, MA 02115

48. Dr. Jim Petty1
 National Biological Survey
 4200 New Haven Road
 Columbia, MO 65201

49. Prof. Maria Rieders.....1
 Dept. of Industrial Eng.
 Northwestern Univ.
 Evanston, IL 60208

50. Dr. John E. Rolph.....1
 RAND Corporation
 1700 Main St.
 Santa Monica, CA 90406

51. Prof. M. Rosenblatt1
Department of Mathematics
University of California, San Diego
La Jolla, CA 92093

52. Prof. Frank Samaniego1
Statistics Department
University of California
Davis, CA 95616

53. Prof. W. R. Schucany1
Dept. of Statistics
Southern Methodist University
Dallas, TX 75222

54. Prof. D. C. Siegmund1
Dept. of Statistics
Sequoia Hall
Stanford University
Stanford, CA 94305

55. Prof. H. Solomon1
Department of Statistics
Sequoia Hall
Stanford University
Stanford, CA 94305

56. Andrew Solow1
Woods Hole Oceanographic Institute
Woods Hole, MA 02543

57. Prof. W. Stuetzle1
Department of Statistics
University of Washington
Seattle, WA 98195

58. Dr. D. J. Svensgaard1
Health Effects Research Lab
U. S. Environmental Protection Agency
Biostatistics Branch MD55
Research Triangle Park, NC 27711

59. Prof. J. W. Tukey1
 Statistics Dept., Fine Hall
 Princeton University
 Princeton, NJ 08540

60. Prof. David L. Wallace1
 Statistics Dept., University of Chicago
 5734 S. University Ave.
 Chicago, IL 60637

61. Dr. Ed Wegman.....1
 George Mason University
 Fairfax, VA 22030

62. Dr. Alan Weiss1
 Rm. 2C-118
 AT&T Bell Laboratories
 600 Mountain Avenue
 Murray Hill, NJ 07974-2040

63. Dr. P. Welch1
 IBM Research Laboratory
 Yorktown Heights, NY 10598

64. Prof. Roy Welsch.....1
 Sloan School
 M.I.T.
 Cambridge, MA 02139